



Received 20 Oct 2016;
Accepted 12 Nov 2016
Online November 2016

G. Vanthana, Research
Scholar, Research
Scholar, Department of
Zoology, T. B.M.L
College, Porayar,
Nagapattinum District,
Tamil Nadu, S. India

*Corresponding author
Dr. AS. Paul Ravindran,
Associate Professor &
Head, Department of
Zoology, T. B.M.L
College, Porayar,
Nagapattinum District
Tamil Nadu, S. India

Research Article

Zoology

**DETERMINATION OF HEAVY METALS AND TRACE
ELEMENTS FROM KUDUVAIYR ESTUARY,
NAGAPATTINUM EAST COAST, NAGAPATTINUM
DISTRICT, TAMIL NADU INDIA**

G. Vanthana# and A.S. Paul Ravindran *

ABSTRACT

A study has been undertaken to analysis of heavy metals and trace elements in Kuduvaier estuary, Nagapattinum east coast, Nagapattinum district, Tamil Nadu India and compared with standard. In the present study we observed that most of the heavy metals and trace elements increased in Kuduvaier estuary as compared to standard as BIS, ICMR and WHO. The problem of pollution is more pronounced in the aquifers of Kuduvaier estuary region. In addition to human activities, the limited availability of potable water resources are being deteriorated by pollutants due to urbanization, increase in population and over exploitation, intense agricultural activities and industrialization. Hence, it needs for an immediate focus on the monitoring of water quality of this region.

Keywords: Kuduvaier estuary, Metals, Trace elements, Nagapattinum, Estuaries.

Citation: G. Vanthana and A.S. Paul Ravindran (2016) Determination of heavy metals and Trace elements from Kuduvaier estuary, Nagapattinum east coast, Nagapattinum district, Tamil Nadu India. Asian Journal of Innovative Research 1(4): 19-22.

INTRODUCTION

The increasing population, urbanization and industrial sources are given rise to environmental stress and pollution all over the world. In fact, most of the developed countries have already realized that human existence on the earth may be endangered if suitable steps are not taken for the abatement; the pollution to the water bodies causes a serious threat to the mankind (Agarwal, 2005). Estuaries are complex and dynamic environmental components which receive large amounts of contaminants from urban and industrial sites (Leight *et al.*, 2005). Industrialization and urbanization of the coastal region often lead to decrease in coastal resource and destruction of natural defense structures (Zhao *et al.*, 2011). Discharge of agricultural wastes, industrial effluents and urban activities is considered to be the primary sources for increasing nutrient load in nearby aquatic water bodies (Kucuksezgin *et al.*, 2006). Eutrophication is of great environmental distress, leading to complicity in the aquatic environment, causing problems such as formation of algal blooms which results reduction in oxygen levels, leads to mortality of

aquatic fauna and flora and eventually loss of biodiversity (Yadav *et al.*, 2007). India is bestowed with long coastline of 8,129 km and of this 6,000 km is rich in estuaries, creeks, brackish water, lagoons and lakes. In the present study we observed that most of the heavy metals and trace elements increased in Kuduvaier estuary as compared to standard.

MATERIALS AND METHODS

Location of sample Collection

The water sample was collected from Kuduvaier estuary, Nagapattinam east coast, Nagapattinam district, Tamil Nadu India.

Heavy Metal analysis of water sample

The water samples were collected in Polyvinyl chloride container and preserved with 6(N) HNO₃ solution to pH < 2, (10 ml/l sample) for estimation of arsenic and others heavy metals such as iron, copper, cadmium, chromium, mercury, zinc and arsenic etc. and Sulphate anion content was determined by using UV- Vis Spectrophotometer (Specord-40, Analytik Jena). The concentrations of metals were determined by using atomic absorption spectroscopy (AAS) and the results were tabulated.

RESULTS AND DISCUSSION

Pollution of the aquatic environment and its effects on the living resources, especially the fishery resources, has assumed considerable interest as well as importance in the recent times. Most of the rivers which discharge large quantities of water into the coastal marine environment are polluted and these pollutants obviously end up in the inshore coastal waters. The vast marine environment has long been used as a site for the disposal of wastes. In some cases the polluted material is discharged directly into the sea and in other cases the pollutant reaches the rivers and estuaries and finally ends up in the sea. Estuaries, the important contributors of fisheries in India, suffer from severe loss of fish production due to increased industrialization and urbanization along the coastal zone by continuous discharge of industrial effluents (Padmini *et al.*, 2007). Under the influence of a variety of inter-related biotic and abiotic structural compounds and intensive chemical, physical and biological processes, estuaries are highly variable systems (Madhupratap, 1987). The human race is under tremendous threat due to undesired changes in the physical, chemical and biological characteristics of air, water and soil. Due to increased human population, industrialization, use of fertilizers and man- made

activities, water is highly polluted with different harmful contaminants (Patil *et al.*, 2012).

The distribution of metals within the aquatic environment is governed by complex processes of material exchange affected by various natural and anthropogenic activities (Ip *et al.*, 2007). Although metals are natural constituents of the earth's crust and are present in all ecosystems, their concentrations have been dramatically increased by human activities (Guerzoni *et al.*, 1984). Since heavy metals are toxic, persistent and non-degradable in the environment, the contamination of sediments by these elements represent the greatest ecological risk to coastal marine ecosystem (Garcia *et al.*, 2008). Such pollution has several distinct biological effects, including diseases in plant and animal species (Lamb *et al.*, 1991), local or complete extinction of some species (Vermeij, 1993) and loss or modification of habitat (Nee and May, 1992).

Heavy metals are persistent pollutants in aquatic ecosystems. The trace metal occurs in all compartments of the marine environment and has a tendency to accumulate in organisms from different trophic levels of marine food webs. The accumulation of trace metals in aquatic organisms can pose a long-term burden on biogeochemical cycling in the ecosphere. Bioaccumulation becomes an environmental problem when chemicals accumulated are toxic. Toxicity may occur along the food chain when the contaminated species or a substance is consumed (Heng *et al.*, 2004). Once trace metals enter the food chain, they may accumulate to dangerous levels and be harmful to human health (Manahan, 2000). Marine fishu have been widely employed as biomonitors for heavy metals pollution due to their ability to accumulate metals without harming themselves. The reliable use of marine molluscs as indicator organisms for metal contamination requires an understanding of how environmental parameters effect metal accumulation. Particularly bivalve molluscs have been considered as a potential biomonitor for metallic contamination in marine ecosystems (Jung and Zauke, 2008). The problem of pollution is more pronounced in the aquifers of Kuduvaier estuary region. In addition to human activities, the limited availability of potable water resources are being deteriorated by pollutants due to urbanization, increase in population and over exploitation, intense agricultural activities and industrialization. Hence, it needs for an immediate focus on the monitoring of water quality of this region.

Table 1 represent the analysis of metals in Kuduvaiyr estuary, Nagapattinum east coast, Nagapattinum district, Tamil Nadu India and compared with standard (BIS, 1998; ICMR, 1994; WHO, 1998). In the present study we observed that most of the heavy metals and trace elements increased in Kuduvaiyr estuary as compared to standard as BIS, ICMR and WHO (Table 2).

Table 1 shows the analysis of metals in Kuduvaiyr estuary, Nagapattinum east coast, Nagapattinum district, Tamil Nadu India

S.No	Parameters	Results
1.	Chloride (mg/l)	878
2.	Sulphate (mg/l)	336
3.	Phosphate (mg/l)	0.20
4.	Nitrate (mg/l)	60
5.	Nitrite (mg/l)	20
6.	Calcium (mg/l)	368
7.	Magnesium (mg/l)	192
8.	Sodium (mg/l)	465
9.	Potassium (mg/l)	1.37
10.	Zinc (mg/l)	12
11.	Copper (mg/l)	2.04
12.	Iron (mg/l)	0.97
13.	Manganese (mg/l)	42
14.	Cadmium (mg/l)	0.12
15.	Selenium (mg/l)	0.006
16.	Arsenic (mg/l)	0.09
17.	Mercury (mg/l)	0.016
18.	Boron (mg/l)	0.08

Table 2 Water quality standards (BIS, ICMR and WHO)

.No.	Parameters	BIS (1998)		ICMR (1994)		WHO (1998)	
		P	E	P	E	P	E
1.	Calcium	75	200	75	200	75	200
2.	Magnesium	30	100	50	150	50	150
3.	Sodium	-	-	-	-	200	-
4.	Potassium	-	-	-	-	-	-
5.	Chloride	250	1000	250	1000	200	600
6.	Sulphate	200	400	200	400	200	400
7.	Nitrate	40	40	20	50	-	50-100
8.	Phosphate	-	-	-	-	-	0.001
9.	Manganese	20	80	30	100	20	100
10.	Cadmium	0.01	0.01	-	0.05	-	0.05
11.	Selenium	--	-	-	--	-	-
12.	Arsenic	0.05	-	-	0.2	-	0.2
13.	Mercury	0.001	0.001	-	-	-	-
14.	Zinc	5	15	-	-	-	-
15.	Copper	0.05	1.5	1.0	3.0	1.0	1.5
16.	Iron	0.3	10.3	1	0.3	1	0.3

Note: P = Permissible limit; E = Excessive limit All parameters are expressed in mg/L except pH, Turbidity (NTU),

CONCLUSION

It is observed from this study, that the maximum concentrations of all the metals in the water are observed in samples collected from Kuduvaivr estuary. It may be due to the discharge of untreated effluent from various industries located near Kuduvaivr. As far as the importance of fish in the human diet is concerned, it is necessary that the biological monitoring of water and fish should be done periodically to ensure the safety of seafood consumption. The safe disposal of industrial effluents and domestic sewage should be practiced to avoid such contamination. Also, the laws enacted to protect the environment should be enforced effectively.

REFERENCES

- Agarwal S., Environ. Poll., APHA Publications, 9th Ed., 23-42 (2005)
- BIS, (1998) specification for drinking water, Bureau of Indian standard, New Delhi. 171.-178.
- Garcia, E.M., Cruz-Motta, J.J., Farina, O., and Bastidas, C. (2008). Anthropogenic influences on heavy metals across marine habitats in the western coast of Venezuela, *Continental Shelf Research*, 28, 2757–2766.
- Guerzoni, S., Frignani, M., Giordani, P., and Frasari, F. (1984). Heavy metals in sediments from different environments of a Northern Adriatic Sea area, Italy, *Environmental Geology and Water Sciences*, 6 (2), 111–119.
- Heng LY, Mokhtar MB, Rusin S (2004) The bioaccumulation of Trace Essential Metals by the freshwater Snail *Turritella* sp. Found in the Rivers of Borneo East Malaysia. *J Bio Sci* 4: 441-444.
- ICMR. (1994). Indian Council of Medical Research. Manual of Standards of Quality of Drinking Water Supplies. 2nd ed. special, report series no. 44, New Delhi
- Ip,C.C.M., Li,X.D., Zhang,G., Wai,O.W.H, and Li,Y.S. (2007). Trace metal distribution in sediments of the Pearl River Estuary and the surrounding coastal areas, South China, *Environmental Pollution*, 147, 311–323.
- Jung K, Zauke GP (2008) Bioaccumulation of trace metals in the brown shrimp *Crangon crangon* (Linnaeus, 1758) from the German Wadden Sea. *Aquat Toxicol* 88: 243-249.
- Kucuksezgin F., Kontas A., Altay O., Uluturhan, E. and Darilmaz E., Assessment of marine pollution in Izmir Bay Nutrient, heavy metal and total hydrocarbon concentrations, *Environ. Int.*, 32, 41-51 (2006)
- Lamb, T., Bickham, J.W., and Gibbons, J.W. (1991). Genetic damage in a population of slider turtles (*Trachemys scripta*) inhabiting a radioactive reservoir, *Archives of Environmental Contamination and Toxicology*, 20, 138-142.
- Leight A.K., Scott G.I., Fulton M.H. and Daugomah J.W., Long term monitoring of grass shrimp *Palaemonetes* sp. Population metrics at sites with agricultural runoff influences, *Integr. Comp. Biol.*, 45, 143-150 (2005)
- Madhuratap M. Status and strategy of zooplankton of tropical Indian estuaries: A Review, *Bulletin of Plankton society of Japan*, 1987; 34(1): 65-81..
- Manahan SE (2000) Environmental Chemistry. Lewis Publishers, Boca Raton, Florida, USA.
- Nee, S., and May, R.M. (1992). Dynamics of meta-populations: habitat destruction and competitive coexistence, *Journal of Animal Ecology*, 61, 37-40.
- Padmini, E, Geetha BV. A comparative seasonal pollution assessment study on Ennore Estuary with respect to metal accumulation in the grey mullet, *Mugil cephalus*. *Int. J. Oceanog. Hydrobiol.*, 2007; 34(4): 91-103.
- Patil, P.N., Sawant, D.V., Deshmukh, R.N. 2012. Physicochemical parameters for testing of water A review. *Int. J. Environ. Sci.*, 3(3): 1194-1207.
- Vermeij, G.J. (1993). Biogeography of recently existing marine species: implications for conservation. *Conservation Biology*, 7, 391-397.
- WHO (1995). Guidelines for drinking water quality. WHO Geneva 2nd Edition (1).
- Yadav A., Gopesh A., Pandey R.S., Rai D.K. and Sharma B., Fertilizer industry effluent induced Biochemical changes in freshwater teleost *Channa striatus* (Bloch), *Bull. Environ. Contam. Toxicol.*, 79, 588-595 (2007)
- Zhao X., Shen Z.Y., Xiong M. and Qi J., Key uncertainty sources analysis of water quality model using the first order error method, *Int. J. Environ. Sci. Tech.*, 8(1), 137-148 (2011)

Source of support: Nil;
Conflict of interest: None declared